

## Levofloxacin 250mg and 500mg Tablets

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- The full name of this medicine is Levofloxacin 250mg and 500mg Tablets but within the leaflet it will be referred to as Levofloxacin tablets.

**What is in this leaflet:**

- 1. What Levofloxacin tablets are and what they are used for**
- 2. What you need to know before you take Levofloxacin tablets**
- 3. How to take Levofloxacin tablets**
- 4. Possible side effects**
- 5. How to store Levofloxacin tablets**
- 6. Contents of the pack and other information**

### **1. What Levofloxacin tablets are and what they are used for**

The name of your medicine is Levofloxacin tablets. Levofloxacin tablets contain a medicine called levofloxacin. This belongs to a group of medicines called antibiotics. Levofloxacin is a "quinolone" antibiotic.

It works by killing the bacteria that cause infections in your body.

Levofloxacin can be used to treat infections of the:

- sinuses
- lungs, in people with long-term breathing problems or pneumonia
- urinary tract, including your kidneys or bladder
- prostate gland, where you have a long lasting infection
- skin and underneath the skin, including muscles. This is sometimes called "soft tissue".

In some special situations, Levofloxacin tablets may be used to lessen the chances of getting a pulmonary disease named anthrax or worsening of the disease after you are exposed to the bacteria causing anthrax.

### **2. What you need to know before you take Levofloxacin tablets**

**Do not take Levofloxacin tablets if you:**

- are **allergic** to levofloxacin, any other quinolone antibiotic such as moxifloxacin, ciprofloxacin or ofloxacin or any of the other ingredients of this medicine (listed in section 6). Signs of an allergic reaction include: a rash, swallowing or breathing problems, swelling of your lips, face, throat or tongue
- have ever had epilepsy
- have ever had a problem with your tendons such as tendonitis that was related to treatment with a 'quinolone antibiotic'. A tendon is the cord that joins your muscle to your skeleton
- are a child or a growing teenager
- are pregnant, might become pregnant or think you may be pregnant

- are breast-feeding

Do not take this medicine if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Levofloxacin tablets.

## **Warnings and precautions**

### **Talk to your doctor or pharmacist before taking Levofloxacin tablets if you:**

- are 60 years of age or older
- are using corticosteroids, sometimes called steroids (see section “Other medicines and Levofloxacin tablets”)
- have ever had a fit (seizure)
- have had damage to your brain due to a stroke or other brain injury
- have kidney problems
- have something known as ‘glucose – 6 – phosphate dehydrogenase deficiency’. You are more likely to have serious problems with your blood when taking this medicine
- have ever had mental health problems
- have ever had heart problems: caution should be taken when using this kind of medicine, if you were born with or have family history of prolonged QT interval (seen on ECG, electrical recording of the heart), have salt imbalance in the blood (especially low level of potassium or magnesium in the blood), have a very slow heart rhythm (called ‘bradycardia’), have a weak heart (heart failure), have a history of heart attack (myocardial infarction), you are female or elderly or you are taking other medicines that result in abnormal ECG changes or are taking vitamin K antagonists (e.g. warfarin), due to a possible increase in coagulation tests and/or bleeding (see section “Other medicines and Levofloxacin tablets”)
- are diabetic
- have ever had liver problems
- have myasthenia gravis
- have been diagnosed with an enlargement or "bulge" of a large blood vessel (aortic aneurysm or large vessel peripheral aneurysm)
- have experienced a previous episode of aortic dissection (a tear in the aorta wall)
- have been diagnosed with leaking heart valves (heart valve regurgitation)
- have a family history of aortic aneurysm or aortic dissection or congenital heart valve disease, or other risk factors or predisposing conditions (e.g. connective tissue disorders such as Marfan syndrome, or vascular Ehlers-Danlos syndrome, Turner syndrome, Sjögren’s syndrome [an inflammatory autoimmune disease], or vascular disorders such as Takayasu arteritis, giant cell arteritis, Behçet’s disease, high blood pressure, or known atherosclerosis, rheumatoid arthritis [a disease of the joints] or endocarditis [an infection of the heart])
- have ever developed a severe skin rash or skin peeling, blistering and/or mouth sores after taking levofloxacin.

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking Levofloxacin tablets.

### **Before taking this medicine**

You should not take fluoroquinolone/quinolone antibacterial medicines, including Levofloxacin tablets, if you have experienced any serious adverse reaction in the past when taking a quinolone or fluoroquinolone. In this situation, you should inform your doctor as soon as possible.

### **When taking this medicine**

- Pain and swelling in the joints and inflammation or rupture of tendons may occur rarely. Your risk is increased if you are elderly (above 60 years of age), have received an organ transplant, have kidney problems or if you are being treated with corticosteroids. Inflammation and ruptures of tendons may occur within the first 48 hours of treatment and even up to several months after stopping of Levofloxacin tablets therapy. At the first sign of pain or inflammation of a tendon (for example in

your ankle, wrist, elbow, shoulder or knee), stop taking Levofloxacin tablets, contact your doctor and rest the painful area. Avoid any unnecessary exercise as this might increase the risk of a tendon rupture.

- You may rarely experience symptoms of nerve damage (neuropathy) such as pain, burning, tingling, numbness and/or weakness especially in the feet and legs or hands and arms. If this happens, stop taking Levofloxacin tablets and inform your doctor immediately in order to prevent the development of a potentially irreversible condition.

If you feel sudden, severe pain in your abdomen, chest or back, which can be symptoms of aortic aneurysm and dissection, go immediately to a hospital accident and emergency department. Your risk may be increased if you are being treated with systemic corticosteroids.

If you start experiencing a rapid onset of shortness of breath, especially when you lie down flat in your bed, or you notice swelling of your ankles, feet or abdomen, or a new onset of heart palpitations (sensation of rapid or irregular heartbeat), you should inform a doctor immediately.

### **Serious skin reactions**

Serious skin reactions including Stevens-Johnson Syndrome (SJS), Toxic Epidermal Necrolysis (TEN) and drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported with the use of levofloxacin.

- SJS/TEN can appear initially as reddish target-like spots or circular patches often with central blisters on the trunk. Also, ulcers of mouth, throat, nose, genitals and eyes (red and swollen eyes) can occur. These serious skin rashes are often preceded by fever and/or flu-like symptoms. The rashes may progress to widespread peeling of the skin and life-threatening complications or be fatal.
- DRESS appears initially as flu-like symptoms and a rash on the face then an extended rash with a high body temperature, increased levels of liver enzymes seen in blood tests and an increase in a type of white blood cell (eosinophilia) and enlarged lymph nodes.

If you develop a serious rash or another of these skin symptoms, stop taking levofloxacin and contact your doctor or seek medical attention immediately.

### **Prolonged, disabling and potentially irreversible serious side effects**

Fluoroquinolone/quinolone antibacterial medicines, including Levofloxacin tablets, have been associated with very rare but serious side effects, some of them being long lasting (continuing months or years), disabling or potentially irreversible. This includes tendon, muscle and joint pain of the upper and lower limbs, difficulty in walking, abnormal sensations such as pins and needles, tingling, tickling, numbness or burning (paraesthesia), sensory disorders including impairment of vision, taste and smell, and hearing, depression, memory impairment, severe fatigue, and severe sleep disorders.

If you experience any of these side effects after taking Levofloxacin tablets, contact your doctor immediately prior to continuing treatment. You and your doctor will decide on continuing the treatment considering also an antibiotic from another class.

### **Children and adolescents**

This medicine must not be given to children or teenagers.

### **Other medicines and Levofloxacin tablets**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This is because Levofloxacin tablets can affect the way some other medicines work. Also some medicines can affect the way Levofloxacin tablets work.

**In particular, tell your doctor if you are taking any of the following medicines. This is because it can increase the chance of you getting side effects, when taken with Levofloxacin tablets:**

- Corticosteroids, sometimes called steroids – used for inflammation. You may be more likely to have

inflammation and/or rupture of your tendons

- Vitamin K antagonists such as Warfarin - used to thin the blood. You may be more likely to have a bleed. Your doctor may need to take regular blood tests to check how well your blood can clot
- Theophylline - used for breathing problems. You are more likely to have a fit (seizure) if taken with Levofloxacin tablets
- Non-steroidal anti-inflammatory drugs (NSAIDs) - used for pain and inflammation such as aspirin, ibuprofen, fenbufen, ketoprofen and indometacin. You are more likely to have a fit (seizure) if taken with Levofloxacin tablets
- Ciclosporin - used after organ transplants. You may be more likely to get the side effects of ciclosporin
- Medicines known to affect the way your heart beats. This includes medicines used for abnormal heart rhythm (antiarrhythmics such as quinidine, hydroquinidine, disopyramide, sotalol, dofetilide, ibutilide and amiodarone), for depression (tricyclic antidepressants such as amitriptyline and imipramine), for psychiatric disorders (antipsychotics), and for bacterial infections ('macrolide' antibiotics such as erythromycin, azithromycin and clarithromycin)
- Probenecid (used for gout) or cimetidine (used for ulcers and heartburn). Special care should be taken when taking either of these medicines with Levofloxacin tablets. If you have kidney problems, your doctor may want to give you a lower dose.

**Do not take Levofloxacin tablets at the same time as the following medicines. This is because it can affect the way Levofloxacin tablets work:**

- Iron tablets (for anaemia), zinc supplements, magnesium or aluminium-containing antacids (for acid or heartburn), didanosine, or sucralfate (for stomach ulcers). See section 3 "If you are already taking iron tablets, zinc supplements, antacids, didanosine or sucralfate" below.

#### **Urine tests for opiates**

Urine tests may show 'false-positive' results for strong painkillers called 'opiates' in people taking Levofloxacin tablets. If your doctor has prescribed a urine test, tell your doctor you are taking Levofloxacin tablets.

#### **Tuberculosis tests**

This medicine may cause "false negative" results for some tests used in laboratory to search for the bacteria causing tuberculosis.

#### **Pregnancy and breast-feeding**

Do not take this medicine if you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby.

#### **Driving and using machines**

You may get side effects after taking this medicine, including feeling dizzy, sleepy, a spinning feeling (vertigo) or changes to your eyesight. Some of these side effects can affect you being able to concentrate and your reaction speed. If this happens, do not drive or carry out any work that requires a high level of attention.

#### **Levofloxacin tablets contain lactose**

If you have been told by your doctor that you have intolerance to some sugars, contact your doctor before taking this medicinal product.

#### **Levofloxacin tablets contain sodium**

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

### **3. How to take Levofloxacin tablets**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

**Taking this medicine**

- Take this medicine by mouth
- Swallow the tablets without crushing with a drink of water
- The tablets can be divided into equal doses
- The tablets may be taken during meals or at any time between meals

**Protect your skin from sunlight**

Keep out of direct sunlight while taking this medicine and for 2 days after you stop taking it. This is because your skin will become much more sensitive to the sun and may burn, tingle or severely blister if you do not take the following precautions:

- Make sure you use high factor sun cream
- Always wear a hat and clothes which cover your arms and legs
- Avoid sun beds

**If you are already taking iron tablets, zinc supplements, antacids, didanosine or sucralfate**

- Do not take these medicines at the same time as Levofloxacin tablets. Take your dose of these medicines at least 2 hours before or after Levofloxacin tablets.

**How much to take**

- Your doctor will decide on how many Levofloxacin tablets you should take
- The dose will depend on the type of infection you have and where the infection is in your body
- The length of your treatment will depend on how serious your infection is
- If you feel the effect of your medicine is too weak or strong, do not change the dose yourself, but ask your doctor.

**Adults and the elderly****Sinuses infection**

- Two tablets of Levofloxacin tablets 250mg, once each day
- Or, one tablet of Levofloxacin tablets 500mg, once each day

**Lung infection, in people with long-term breathing problems**

- Two tablets of Levofloxacin tablets 250mg, once each day
- Or, one tablet of Levofloxacin tablets 500mg, once each day

**Pneumonia**

- Two tablets of Levofloxacin tablets 250mg, once or twice each day
- Or, one tablet of Levofloxacin tablets 500mg, once or twice each day

**Infection of urinary tract, including your kidneys or bladder**

- One or two tablets of Levofloxacin tablets 250mg, each day
- Or, 1/2 or one tablet of Levofloxacin tablets 500mg, each day

**Prostate gland infection**

- Two tablets of Levofloxacin tablets 250mg, once each day
- Or, one tablet of Levofloxacin tablets 500mg, once each day

**Infection of skin and underneath the skin, including muscles**

- Two tablets of Levofloxacin tablets 250mg, once or twice each day
- Or, one tablet of Levofloxacin tablets 500mg, once or twice each day

**Inhalation anthrax exposure**

- Two tablets of Levofloxacin tablets 250mg, once each day

- Or, one tablet of Levofloxacin tablets 500mg, once each day

#### **Adults and the elderly with kidney problems**

Your doctor may need to give you a lower dose.

#### **Use in children and adolescents**

This medicine must not be given to children or teenagers.

#### **If you take more Levofloxacin tablets than you should**

If you accidentally take more tablets than you should, tell a doctor or get other medical advice straight away. Take the medicine pack with you. This is so the doctor knows what you have taken. The following effects may happen: convulsive fits (seizures), feeling confused, dizzy, less conscious, having tremor and heart problems - leading to uneven heartbeats as well as feeling sick (nausea) or having stomach burning.

#### **If you forget to take Levofloxacin tablets**

If you forgot to take a dose, take it as soon as you remember unless it is nearly time for your next dose. Do not take a double dose to make up for a forgotten dose.

#### **If you stop taking Levofloxacin tablets**

Do not stop taking Levofloxacin Tablets just because you feel better. It is important that you complete the course of tablets that your doctor has prescribed for you. If you stop taking the tablets too soon, the infection may return, your condition may get worse or the bacteria may become resistant to the medicine.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them. These effects are normally mild or moderate and often disappear after a short time.

**Stop taking Levofloxacin tablets and see a doctor or go to a hospital straight away if you notice the following side effects:**

**Not known (cannot be estimated from available data)**

- You have an allergic reaction. The signs may include: a rash, swallowing or breathing problems, swelling of your lips, face, throat, or tongue

**Stop taking Levofloxacin tablets and see a doctor straight away if you notice any of the following serious side effects - you may need urgent medical treatment:**

**Rare (may affect up to 1 in 1,000 people)**

- Fits (convulsions)
- Widespread rash, high body temperature, liver enzyme elevations, blood abnormalities (eosinophilia), enlarged lymph nodes and other body organs involvement (Drug Reaction with Eosinophilia and Systemic Symptoms which is also known as DRESS or drug hypersensitivity syndrome). See also section 2
- Syndrome associated with impaired water excretion and low levels of sodium (SIADH)
- Bruising and bleeding easily due to a lowering in the number of blood platelets (thrombocytopenia)
- Low number of white blood cells (neutropenia)
- Exaggerated immune response (hypersensitivity)
- Seeing or hearing things that are not there (hallucinations, paranoia), change in your opinion and thoughts (psychotic reactions).

**Not known (cannot be estimated from available data)**

- Serious skin rashes including Stevens-Johnson syndrome and toxic epidermal necrolysis. These can appear as reddish target-like macules or circular patches often with central blisters on the trunk, skin peeling, ulcers of mouth, throat, nose, genitals and eyes and can be preceded by fever and flu-like symptoms. See also section 2
- Loss of appetite, skin and eyes becoming yellow in colour, dark-coloured urine, itching, or tender stomach (abdomen). These may be signs of liver problems which may include a fatal failure of the liver
- Burning, tingling, pain or numbness. These may be signs of something called 'neuropathy'
- Watery diarrhoea which may have blood in it, possibly with stomach cramps and a high temperature. These could be signs of a severe bowel problem
- Pain and inflammation in your tendons or ligaments, which could lead to rupture. The Achilles tendon is affected most often
- Fever, sore throat and a general feeling of being unwell that does not go away. This may be due to a lowering in the number of white blood cells (agranulocytosis)
- Loss of circulation (anaphylactic-like shock)
- Temporary loss of consciousness or posture (syncope)
- Temporary loss of vision
- Abnormal fast heart rhythm, life-threatening irregular heart rhythm including cardiac arrest, alteration of the heart rhythm (called 'prolongation of QT interval', seen on ECG, electrical activity of the heart)
- Pancreatitis
- Inflammation of the liver (hepatitis)
- Muscle rupture and muscle destruction (rhabdomyolysis)
- Risk of having suicidal thoughts or actions
- Attacks of porphyria in people who already have porphyria (a very rare metabolic disease).

If your eyesight becomes impaired or if you have any other eye disturbances whilst taking Levofloxacin tablets, consult an eye specialist immediately.

**Tell your doctor if any of the following side effects get serious or last longer than a few days:**

**Common (may affect up to 1 in 10 people)**

- Sleeping problems
- Headache, feeling dizzy
- Feeling sick (nausea, vomiting) and diarrhoea
- Increase in the level of some liver enzymes in your blood

**Uncommon (may affect up to 1 in 100 people)**

- Changes in the number of other bacteria or fungi, infection by fungi named Candida, which may need to be treated
- Changes in the number of white blood cells shown up in the results of some blood tests (leukopenia, eosinophilia)
- Feeling stressed (anxiety), feeling confused, feeling nervous, feeling sleepy, trembling, a spinning feeling (vertigo)
- Shortness of breath (dyspnoea)
- Changes in the way things taste, loss of appetite, stomach upset or indigestion (dyspepsia), pain in your stomach area, feeling bloated (flatulence) or constipation
- Itching and skin rash, severe itching or hives (urticaria), sweating too much (hyperhidrosis)
- Joint pain or muscle pain
- Blood tests may show unusual results due to liver (bilirubin increased) or kidney (creatinine increased) problems
- General weakness

**Rare (may affect up to 1 in 1,000 people)**

- Lowering of your blood sugar levels (hypoglycaemia). This is important for people that have diabetes

- Feeling depressed, mental problems, feeling restless (agitation), abnormal dreams or nightmares
- Tingly feeling in your hands and feet (paraesthesia)
- Problems with your hearing (tinnitus) or eyesight (blurred vision)
- Unusual fast beating of your heart (tachycardia), awareness of your heartbeat (palpitation) or low blood pressure (hypotension)
- Muscle weakness. This is important in people with myasthenia gravis (a rare disease of the nervous system)
- Changes in the way your kidney works and occasional kidney failure which may be due to an allergic kidney reaction called interstitial nephritis
- Fever
- Sharply demarcated, erythematous patches with/without blistering that develop within hours of administration of levofloxacin and heals with post-inflammatory residual hyperpigmentation; it usually recurs at the same site of the skin or mucous membrane upon subsequent exposure to levofloxacin.

**Other side effects include:**

- Lowering in red blood cells (anaemia): this can make the skin pale or yellow due to damage of the red blood cells; lowering in the number of all types of blood cells (pancytopenia)
- Increase of your blood sugar levels (hyperglycaemia) or lowering of your blood sugar levels leading to coma (hypoglycaemic coma). This is important for people that have diabetes
- Changes in the way things smell, loss of smell or taste (parosmia, anosmia, ageusia)
- Problems moving and walking (dyskinesia, extrapyramidal disorders)
- Impairment or loss of hearing
- Difficulty breathing or wheezing (bronchospasm)
- Allergic lung reactions
- Increased sensitivity of your skin to sun and ultraviolet light (photosensitivity)
- Inflammation of the vessels that carry blood around your body due to an allergic reaction (vasculitis)
- Inflammation of the tissue inside the mouth (stomatitis)
- Joint redness and swelling (arthritis)
- Pain, including pain in the back, chest and extremities
- Persistent headache with or without blurred vision (benign intracranial hypertension).

Very rare cases of long lasting (up to months or years) or permanent adverse drug reactions, such as tendon inflammations, tendon rupture, joint pain, pain in the limbs, difficulty in walking, abnormal sensations such as pins and needles, tingling, tickling, burning, numbness or pain (neuropathy), depression, fatigue, sleep disorders, memory impairment, as well as impairment of hearing, vision and taste and smell have been associated with administration of quinolone and fluoroquinolone antibiotics, in some cases irrespective of pre-existing risk factors.

Cases of an enlargement and weakening of the aortic wall or a tear in the aortic wall (aneurysms and dissections), which may rupture and may be fatal, and of leaking heart valves have been reported in patients receiving fluoroquinolones/quinolones. See also section 2.

**Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Levofloxacin Tablets**

Keep this medicine out of the sight and reach of children.



Do not use this medicine after the expiry date which is stated on the carton and foil after EXP. The expiry date refers to the last day of the month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Levofloxacin tablets contain**

- The active substance is levofloxacin.

Each 250mg film-coated tablet contains 250mg of levofloxacin equivalent to 256.23mg of levofloxacin hemihydrate.

Each 500mg film-coated tablet contains 500mg of levofloxacin equivalent to 512.46mg of levofloxacin hemihydrate.

- The other ingredients are:

For the tablet core: sodium stearyl fumarate, crospovidone, silica colloidal anhydrous, copovidone, microcrystalline cellulose silicified (98% microcrystalline cellulose and 2% silica colloidal)

For the tablet coating: Opadry II Pink (Lactose monohydrate, hypromellose 15 cP, titanium dioxide (E171), triacetin, iron oxide red (E172), iron oxide yellow (E172)).

### **What Levofloxacin tablets look like and contents of the pack**

Levofloxacin 250mg film-coated tablets:

Pink biconvex tablets, scored on one side and marked with "L" on the other side. Approximately 13mm long and 6mm wide. The tablet can be divided into equal doses.

Levofloxacin 500mg film-coated tablets:

Pink biconvex tablets, scored on one side and marked with "L" on the other side. Approximately 16mm long and 8mm wide. The tablet can be divided into equal doses.

Pack sizes: 5 and 10 tablets

### **Marketing Authorisation Holder**

Accord, Barnstaple, EX32 8NS, UK

### **Manufacturer**

Actavis Ltd., BLB015-016 Bulebel Industrial Estate, Zejtun ZTN 3000, Malta

This leaflet was last revised in March 2022.



**If you would like a leaflet with larger text, please contact 01271 385257.**